



Autism Practice Brief

Autism Q&A:

Introduction to Teaching Young Children with Autism Spectrum Disorder to Communicate Using Spoken Language

Many young children with Autism Spectrum Disorder (ASD) have difficulty learning to use spoken language. Some children may be able to use only single words while others may be able to make sounds. Still others might not speak at all. Communication is an essential life skill that leads to enhanced interactions and improved quality of life. For all young children who have communication impairments, building skills is critical. There are many different modes or ways we can teach a child with ASD to communicate. For example, a child can learn to use sign language, communicate by exchanging objects or pictures, or can use an electronic device with voice output. While all of these are effective and valuable modes to communicate, we will also want to focus on teaching the child to communicate using spoken language.

In this Q&A, we will explore how to teach spoken communication to children with ASD. Each child is different and possesses various strengths and skills. Spoken communication may be easier to learn for some children than it is for others. Because of this, spoken communication may be the primary way to communicate for some children, while others might use it to supplement other modes. Regardless, the steps to teaching spoken communication are similar for all young children and can be applied to those on the spectrum who are learning to communicate.

QUESTION: How do caregivers and educators begin teaching spoken communication?

ANSWER

When teaching spoken communication the most important factor will be motivation! Before you start any com-

munication intervention, you must know the child and find things that motivate him or her. What are the child's favorite things to do? What are his or her favorite games, places, people? What makes the child laugh? As you determine the items that motivate the child, consider the following areas: physical activities (running, jumping), toys (trains, dolls), edibles (food), liquids (drinks), activities (computer games), social interactions (playing with a sibling, tickle games, peek-a-boo). It will also be helpful to identify items that are considered to be repetitive or restricted patterns of behavior in which the child engages and enjoys. For example, a child might like watching items spin, enjoy flicking a string up and down, or like driving a car on a track over and over.

QUESTION: When should caregivers and educators teach spoken communication?

ANSWER

Adults should teach when the child is motivated to communicate! This will likely be when the child is engaged in his or her favorite activities. Caregivers and educators can embed strategies into the child's favorite activities and use the child's favorite toys and foods to elicit communication. For example, if the child loves gross motor activities (running, jumping on the bed, being spun and thrown in the air,) then these times can be used to teach him/her to communicate.

When teaching spoken communication, begin by teaching the child to request an item. Requesting is a basic function of communication and the best starting point, as it helps the child to get needs and desires met. Using motivational moments will allow the caregiver or educator to capitalize on the child's desires. If the child is whining to get something or reaching for a desired item, this is the perfect opportunity

QUESTION: What are the steps to teach communication?

ANSWER


1. Interact with the child during the activity. Play with the child and make the activity fun and engaging.
2. Elaborate on the child's communication by providing a model. Label or say the word the child is to learn. If he or she is playing with a ball label it by saying, "ball." As the word is modeled, start by saying the word multiple times, pausing in between and saying the word calmly.
3. Expand the child's language. Eventually pair words together and increase what the child is to say. Here is an example of expanding: "Yellow ball!" "What do you want?" "Throw yellow ball!"
4. Be patient and pause to allow the child to communicate. Provide a model and then wait several seconds. It may be necessary to wait 15 or more seconds to allow the child to process. After multiple language models, begin adding in a pause to allow the child to interact. The goal is to give the child an opportunity to respond and to speak. For example, you can say "What are you playing with? (pause) It's a (pause) ball!"

QUESTION: How can caregivers and parents prompt communication?

ANSWER

When teaching spoken communication, a prompt can be provided. The prompt can provide the entire word or phrase the child is to say. For example, when showing the child a duck the prompt simply would be, "duck." A prompt also can provide only part of the word or phrase. For example, as the child is learning to say the word duck, the prompt may be, "du." This type of prompt provides just enough of a cue to remind the child what he is to say.

When using prompts be sure to fade the prompts or increase the expectation for communication. The ultimate goal should be for the child to communicate independently; however, if the child is constantly being prompted he will not get the opportunity to be an independent communicator.



QUESTION: What are the steps to teach communication?

ANSWER

Shaping is the process in which a caregiver or educator gradually shapes the child's response by slowly accepting closer and closer approximations to the word the child is to say. Below is an example of how communication can be shaped. Here, the caregiver or educator is reinforcing the child in building and shaping the word "bubble."

- A: "Look.--Bubble." (then blow bubbles)
- C: "Aaaah"
- A: "bubble" (then blow bubbles)
- C: "ah ah"
- A: "Yes, bubble" (then blow bubbles)
- C: "bu"
- A: "bubble!" (blow bubbles)
- C: "bu-bu"
- A: "bubbles!" (blow bubbles)

QUESTION: What are the most important tips to remember when teaching a child to communicate using spoken language?

ANSWER

- Teach words that are meaningful and are of interest to the child. Some early words are: all done, up, want, drink, or a favorite food item. Many caregivers teach the words 'please' or 'more' first, which are great words to use for requesting; however their meaning can vary depending on the context. For example, if the child is eating dinner and asks for 'more' it is not clear if they are referring to 'more food' or 'more drink.' By teaching the child the word 'milk' it is obvious what the child wants. Gross motor skills are often a strength for children with ASD, thus teaching verbs first is also very motivating and practical for children.
- Identify the activities of the day when you are going to teach communication. These are the 'additional opportunities' or planned opportunities discussed earlier in this fact sheet. It is best to identify the specific activities so they are remembered and the child is provided ample opportunities to communicate each day.
- Be prepared for multiple trials. Children with ASD may require many learning trials to learn to speak. Every child with ASD is different. Some will learn to imitate and

respond to your vocalizations quickly and some will take longer to respond.

- Incorporate high intensity reactions! These reactions motivate children to speak by getting their attention and getting them excited about the activity. If the caregiver is excited about the activity, the child will be too. Have fun during play!
- Talk! Even if the child is unresponsive. Talk to the child about their play using short sentences. Narrate what they are doing and then pause to let them speak. Caregivers often stop speaking with their child because the child is not responding. Even if the child is unresponsive, never stop talking! Talking helps them process and attend to language, and understand that it is meaningful.
- Praise and reinforce the approximations. Reinforcement for spoken communication should be easy. The child gets the item he or she wants! We have to start somewhere, so praise and reinforce the approximations and the attempts to communicate. Reinforce the child for trying. Do not just reinforce if the word is spoken perfectly.
- Pause to give the child a chance to respond. For example, when a child is pointing to a ball, say “Do you want your ball?” then wait for the child to respond. Children with ASD might take many seconds to process information and respond, so be patient.
- Imitate and elaborate the child’s vocalizations. Imitate and elaborate the vocalizations so they know what they said is meaningful! When the child points at a cookie and says “cookie” you can say, “Oh, you want a COOKIE?” When elaborating and expanding on the vocalizations, it helps them learn to expand their own language.

SUMMARY

Many young children with ASD have difficulty learning to use spoken language. Communication is a critical life skill and many children with ASD can learn to speak. Teaching these children takes time and lots of repetition, but it should be a fun process for the child and caregiver. Communication should build on the child’s motivators and incorporated into daily routines. Prompting can be used to shape the child’s use of words so he or she is able to move from making

from making sounds or approximations to saying words and even phrases.

ADDITIONAL RESOURCES

Additional Q&A Fact Sheets provides information on how to support individuals with ASD in the classroom and community.

Please visit the ACE website for more resources
<http://www.vcuautismcenter.org>

REFERENCES

- Autism Speaks. Symptoms. Retrieved from <http://www.autismspeaks.org/what-autism/symptoms> on 2011, September 19.
- Koegel, L. (2000). Interventions to facilitate communication in autism. *Journal of Autism and Developmental Disorders*, 30(5), 383-391.
- Landa, R. (2007). Early communication development and intervention for children with autism. *Mental Retardation and Developmental Disabilities Research Reviews*, 13, 16-25.
- Levy, S., Kim, A., & Olive, M. (2006). Interventions for young children with autism: A synthesis of the literature. *Focus on Autism and Other Developmental Disabilities*, 21(1), 55-62.
- Paparella, T., Goods, K., Freeman, S., & Kasari, C. (2011). The emergence of nonverbal joint attention and requesting skills in young children with autism. *Journal of Communication Disorders*, 44, 569-583.
- Siller, M. and Sigman, M. (2002). The behaviors of parents of children with autism predict the subsequent development of their children’s communication. *Journal of Autism and Developmental Disorders*, 32 (2), 77-89.
- Whalen, C., Schreibman, L., & Ingersoll, B. (2006) The collateral effects of joint attention training on social initiations, positive affect, imitation, and spontaneous speech for young children with autism. *Journal of Autism Developmental Disorder*, 36, 655-664.

Contributors for this issue:
Heather Fleming, M.Ed. and Dawn Hendricks, Ph.D.

Editor: Becky Boswell, M.B.A.

Information for this Autism Practice Brief is from Virginia Commonwealth University’s Autism Center for Excellence (VCU-ACE), which is funded by the Virginia State Department of Education (Grant # 881-61172-H027A100107).

Virginia Commonwealth University, School of Education and Department of Physical Medicine and Rehabilitation is an equal opportunity/ affirmative action institution providing access to education and employment without regard to age, race, color, national origin, gender, religion, sexual orientation, veteran’s status, political affiliation, or disability. If special accommodations are needed, please contact Carol Schall at (804) 828-1851 VOICE or (804) 828-2494 TTY.